

RELAXATION TIME OF L -REVERSAL CHAINS AND OTHER CHROMOSOME SHUFFLES

N. CANCRINI, P. CAPUTO, AND F. MARTINELLI

ABSTRACT. We prove tight bounds on the relaxation time of the so called L -reversal chain, introduced by R. Durrett as a stochastic model for the evolution of chromosome chains. The process is described as follows: we have n distinct letters on the vertices of the n -cycle ($\mathbb{Z} \bmod n$); at each step a connected subset of the graph is chosen uniformly at random among all those of length at most L and the current permutation is shuffled by reversing the order of the letters over that subset. We show that the relaxation time $\tau(n, L)$, defined as the inverse of the spectral gap of the associated Markov generator, satisfies $\tau(n, L) = O(n \vee \frac{n^3}{L^3})$. Our results can be interpreted as a strong evidence for a conjecture of R. Durrett of a similar behavior for the mixing time of the chain.

1. INTRODUCTION, MODELS AND RESULTS

In a series of recent papers R. Durrett has proposed stochastic models based on shuffling rules for the analysis of the evolution of chromosomes [6, 7, 8]. One of the main issues is that of determining the time needed for such processes to reach stationarity. Besides proving several results in this direction and discussing possible applications he pointed out a number of interesting conjectures. One of them refers to the so-called L -reversal model which is described as follows.

We consider a chain (the chromosome) of n distinct genes, so that the configuration of the system at any given time is a permutation η of n letters over n vertices. The vertices are lying on a circle and we take the graph structure of the n -cycle (where each vertex is connected to exactly two vertices). Given an integer $L \leq n$, one step of the L -reversal process is described as follows: uniformly at random we pick a vertex x and, independently, choose a number $1 \leq \ell \leq L$. Given x and ℓ we perform the transition $\eta \rightarrow \eta^{x,\ell}$, where $\eta^{x,\ell}$ is obtained from η by reversing the order of the letters in the segment $\{x, x+1, \dots, x+\ell\}$ with the sums taken modulo n . Note that the probability for the transition $\eta \rightarrow \eta^{x,\ell}$ coincides with that of the transition $\eta^{x,\ell} \rightarrow \eta$ and therefore we have reversibility with respect to the uniform distribution ν over all $n!$ configurations. As in [6] we consider the continuous time version of the L -reversal dynamics in which the steps described above are performed at the arrival times of an independent rate 1 Poisson process. This defines a continuous time Markov chain converging to the uniform distribution ν .

Date: 6 Dec. 2004.

Key words: L -reversal chains; random transpositions; spectral gap.

2000 MSC: 65C05, 92D10.

Let $T(n, L)$ denote the *mixing time* of the chain, i.e.

$$T(n, L) = \inf \{t > 0 : \|p_t - \nu\|_{TV} \leq 1/e\}, \quad (1.1)$$

where $\|\cdot\|_{TV}$ stands for the total variation distance and p_t denotes the distribution of the configuration at time t for the process started from some arbitrary (i.e. completely ordered) configuration η (note that $\|p_t - \nu\|_{TV}$ does not depend on the chosen η because ν is the uniform measure).

The striking conjecture in [6] (see also Problem 4.1 in [8]) is that the mixing time $T(n, L)$ of the L -reversal process should satisfy

$$\frac{1}{C} (n \vee \frac{n^3}{L^3}) \log n \leq T(n, L) \leq C (n \vee \frac{n^3}{L^3}) \log n, \quad (1.2)$$

for some universal constant $C < \infty$. Note that when $L = 1$ we have the usual transposition dynamics on the n -cycle which has been studied in great detail, see e.g. [4, 5, 12, 10]. In particular the conjecture is known to hold in this case. Of most interest are the cases $L = n^\alpha$, $\alpha \in (0, 1)$.

To support the validity of (1.2) Durrett uses comparison arguments and an adaptation of Wilson's method to estimate mixing times [13]. In this way, by a careful choice of the slow modes of relaxation he obtains the lower estimates in (1.2). As for the upper estimates, by comparison with a suitable random transposition dynamics he proves that

$$T(n, L) \leq C \frac{n^3}{L^2} \log n. \quad (1.3)$$

When $L = n^\alpha$, $\alpha \in (0, 1)$, with respect to (1.2) the estimate (1.3) is off by a factor of order n^α for $\alpha \leq 2/3$ and $n^{2(1-\alpha)}$ for $\alpha \geq 2/3$. This shows that the comparison with transpositions is sharp only in the two extreme regimes $\alpha \in \{0, 1\}$. We are not able to prove (nor disprove) the desired upper bound in (1.2) but we considered the problem of establishing similar estimates for the *relaxation time* $\tau(n, L)$, i.e. the inverse of the spectral gap of the chain. In particular, what we find confirms, at least at the level of the spectral gap, the predicted departure from the random transposition behavior in the intermediate regimes.

To describe our main result we introduce some notations. For any function f of the configurations we write $\nu[f] = \frac{1}{n!} \sum_\eta f(\eta)$ for its expectation and $\text{Var}(f) = \nu[f^2] - \nu[f]^2$ for its variance w.r.t. the uniform measure ν . We also write $f^{x,\ell}$ for the function $\eta \rightarrow f(\eta^{x,\ell})$ and $R_{x,\ell} f = f^{x,\ell} - f$ for the associated gradient. The Dirichlet form of the L -reversal dynamics introduced above is

$$\mathcal{E}(f, f) = \frac{1}{2} \frac{1}{nL} \sum_{x=1}^n \sum_{\ell=1}^L \nu[(R_{x,\ell} f)^2]. \quad (1.4)$$

The relaxation time is then given by

$$\tau(n, L) = \sup_f \frac{\text{Var}(f)}{\mathcal{E}(f, f)}, \quad (1.5)$$

with the supremum taken over all non-constant functions f .

Our main result reads as follows.

Theorem 1.1. *There exists $C < \infty$ such that for any $n \geq 2$, any $1 \leq L \leq n$*

$$\frac{1}{C} (n \vee \frac{n^3}{L^3}) \leq \tau(n, L) \leq C (n \vee \frac{n^3}{L^3}). \quad (1.6)$$

As usual the lower estimate in (1.6) can be obtained by a suitable choice of test functions. To obtain a sharp upper bound on $\tau(n, L)$ the argument is more delicate. Here we combine new comparison arguments with an improved iteration scheme inspired by recent work of Carlen, Carvalho and Loss [3]. Our method allows to derive sharp estimates for more general models in the class of so-called p -reversal processes, where each inversion $\eta \rightarrow \eta^{x,\ell}$ is performed with probability $p(\ell)$, with p a probability vector on $\ell \in \{1, \dots, n\}$. The L -reversal process (1.4) is then obtained for $p(\ell) = \frac{1}{L}$, $\ell = 1, \dots, L$. The so-called θ -reversal process is defined by $p(\ell) \propto \theta^\ell$, $\theta \in (0, 1)$, with Dirichlet form

$$\mathcal{E}_\theta(f, f) = \frac{1}{2} \frac{(1-\theta)}{n} \sum_{x=1}^n \sum_{\ell=1}^n \theta^{\ell-1} \nu[(R_{x,\ell}f)^2]. \quad (1.7)$$

In this case, calling $\tau(n, \theta)$ the relaxation time associated to (1.7) we obtain

Theorem 1.2. *There exists $C < \infty$ such that for any $n \geq 2$, any $\theta \in (0, 1)$*

$$\frac{1}{C} (n \vee [n(1-\theta)]^3) \leq \tau(n, \theta) \leq C (n \vee [n(1-\theta)]^3). \quad (1.8)$$

We conclude this introduction with a last remark on Durrett's conjecture (1.2). It is well known that $T(n, L) \geq \tau(n, L)/C$. On the other side an application of standard bounds (see e.g. Theorem 2.1.7 in [12]) gives only that $T(n, L) \leq C \tau(n, L) \log(n!)$. Better estimates can be obtained if one considers the decay of relative entropy functionals rather than L^2 -norms. Namely, let $\beta(n, L)$ denote the *entropy dissipation constant* given by

$$\beta(n, L) = \sup_{f>0} \frac{\text{Ent}(f)}{\mathcal{E}(f, \log f)}, \quad (1.9)$$

with the supremum taken over all positive non-constant functions and we use the notation $\text{Ent}(f) = \nu[f \log f] - \nu[f] \log \nu[f]$ for the entropy of f . It is not hard (see e.g. Corollary 2.8 in [1]) to obtain the bound

$$T(n, L) \leq C \beta(n, L) \log \log(n!).$$

Therefore the conjecture (1.2) would follow if we could prove the bounds (1.6) with τ replaced by β . We believe it is a challenging open question whether our analysis can be extended to treat also the entropy dissipation constant. It is perhaps interesting to observe that the predicted mixing time cannot be derived from logarithmic Sobolev inequalities here. Indeed, we will show that if $s(n, L)$ denotes the log-Sobolev constant (obtained as in (1.9) with $\mathcal{E}(\sqrt{f}, \sqrt{f})$ in place of $\mathcal{E}(f, \log f)$) then

$$s(n, L) \geq \frac{n^2}{C L}. \quad (1.10)$$

With respect to the $\tau(n, L)$ of Theorem 1.1, for $L = n^\alpha$, the bound (1.10) shows that $s(n, L) \geq \tau(n, L) n^{2\alpha-1}$, for $1/2 \leq \alpha \leq 2/3$ and $s(n, L) \geq \tau(n, L) n^{1-\alpha}$ for $\alpha \geq 2/3$. While comparison with random transposition can be always used to prove an upper bound of the form $s(n, L) \leq C(n^3/L^2) \log n$ (see e.g. Theorem 6 in [6]), it remains an open question to understand the true asymptotic behavior in L, n for the log-Sobolev constant.

The rest of the paper consists of two sections: in the first we shortly outline the proof of the upper bound on the relaxation time, develop all the technical parts of our approach and finally prove the corresponding estimates in Theorem 1.1 and Theorem 1.2. In the final section we prove the matching lower bounds and derive the estimate (1.10) on the log-Sobolev constant.

2. UPPER BOUNDS

The main idea for the proof of the upper bounds in Theorems 1.1, 1.2 is to compare the L -reversal chain with (an average over $\ell \leq L$ of) a suitable “block dynamics” on scale ℓ that we call the *long-range average-exchange block dynamics*. Such an auxiliary process seems to be at the heart of several shuffling chains and it can be described informally as follows (see the subsection (2.3) below).

Assume for simplicity that $n = N\ell$ for some integer N and partition the vertices $\{1, 2, \dots, n\}$ into N non-overlapping blocks $\{I_i\}$ of length ℓ . Then, with rate $1/N$, a pair of blocks is chosen, say I_i and I_j , and, by flipping a fair coin, one of the following operations is performed:

- (a) the configuration of letters inside I_i is exchanged with that inside I_j ;
- (b) the letters inside $I_i \cup I_j$ are uniformly re-shuffled over the vertices of $I_i \cup I_j$.

A first key step consists then in proving that the relaxation time of the above block dynamics is $O(1)$ uniformly in N, ℓ (see Proposition 2.3). This step is carried out via an adaptation and extension of the techniques introduced by Carlen, Carvalho and Loss [3] for the Kac model.

The comparison of the long-range average-exchange block dynamics to the L -reversal chain is now done in two further steps.

In the first step we compare the long-range average-exchange block dynamics to an intermediate block dynamics in which moves of type (a) or (b) are performed only between *neighboring blocks* with moves of type (a) occurring at rate N^2 and moves of type (b) occurring at rate 1. We may call this last process the *local average-exchange block dynamics*. This comparison uses rather standard path techniques but the crucial point is that the usual diffusive scaling factor N^2 appears only in front of the exchange operations and does not multiply the local averaging moves (see Lemma 2.5).

In the second step we compare the local average-exchange block dynamics to the L -reversal chain. This presents little difficulty. In fact the contribution of local exchanges (moves of type (a)) is easily expressed in terms of (three) suitable ℓ -reversal moves $R_{x,\ell}$ (see Lemma 2.6). Moreover, the contribution coming from the local averaging (moves of type (b)) on a pair of neighboring blocks I_i and I_{i+1} ,

can be related to inversions. This uses Poincaré inequality for the usual transposition chain restricted to $I_i \cup I_{i+1}$ and a straightforward comparison between the latter and a 2ℓ -reversal chain on $I_i \cup I_{i+1}$ (see Lemma 2.7). The key observation here is that this last step does not spoil the estimates because, as we just learned, transposition and L -reversal chains can be successfully compared if $L = n$.

2.1. Setting and notation. The setting and frequently used notations are described as follows. V_n is the set of n ordered vertices $\{1, 2, \dots, n\}$. Unless otherwise stated the sum $x + y$ is assumed to be taken modulo n , for any $x, y \in V_n$. A permutation (often called configuration) of n letters over V_n is denoted by η , with $\eta_x = j$ meaning that we have letter j at vertex $x \in V_n$. We write Ω for the space of all $n!$ configurations. ν is the uniform distribution on Ω . For any integer $m \leq n$, $I \subset V_n$ is called an m -block if I is of the form $\{x + 1, x + 2, \dots, x + m\}$ for some $x \in V_n$. Given $k \in \mathbb{N}$ we write $\eta_{k,m}$ for the configuration over the m -block $\{(k-1)m+1, (k-1)m+2, \dots, km\}$, i.e. $\eta_{1,m}$ gives the letters over the first m -block, $\eta_{2,m}$ those over the second m -block and so on. Let also Ω_m denote the set of possible configurations over the the first m -block (i.e. all possible realizations of $\eta_{1,m}$).

2.2. Preliminary spectral estimates. We start with a spectral computation that will be used in the arguments below. For any $m \leq n$, let \mathcal{K} denote the symmetric stochastic matrix given by the conditional probabilities

$$\mathcal{K}(\alpha, \beta) := \nu[\eta_{1,m} = \beta \mid \eta_{2,m} = \alpha], \quad \alpha, \beta \in \Omega_m. \quad (2.1)$$

Lemma 2.1. *For any $m \leq n/2$, the spectrum of \mathcal{K} is given by*

$$\lambda_k := (-1)^k \binom{n-m-k}{m-k} \binom{n-m}{m}^{-1}, \quad k = 0, \dots, m \quad (2.2)$$

Proof. Let \mathcal{H}_m denote the space of all functions $\varphi : \Omega_m \rightarrow \mathbb{R}$ and write

$$\langle \varphi, \psi \rangle = \frac{1}{n!} \sum_{\eta \in \Omega} \varphi(\eta_{1,m}) \psi(\eta_{1,m}) = \frac{(n-m)!}{n!} \sum_{\alpha \in \Omega_m} \varphi(\alpha) \psi(\alpha) \quad (2.3)$$

for the scalar product of $\varphi, \psi \in \mathcal{H}_m$. We write $\mathcal{K}\varphi$ for the function

$$\mathcal{K}\varphi(\alpha) = \sum_{\beta \in \Omega_m} \mathcal{K}(\alpha, \beta) \varphi(\beta).$$

By definition of $\mathcal{K}(\alpha, \beta)$, $\mathcal{K}\varphi$ does not depend on the order of the m letters, for any $\varphi \in \mathcal{H}_m$. If $\bar{\mathcal{H}}_m$ denotes the linear subspace of functions $\varphi \in \mathcal{H}_m$ that are independent of the order of the m letters, then any eigenvector of \mathcal{K} must be in $\bar{\mathcal{H}}_m$. Clearly, constant functions give the eigenvalue $\lambda_0 = 1$.

We say that letter j belongs to $\alpha \in \Omega_m$ and write $j \in \alpha$ if the letter labeled j appears in the configuration α . For any $j = 1, \dots, n$ we define χ_j as the indicator function of the event $\{j \in \alpha\}$. To determine the spectrum we may proceed as follows. Consider $1 \leq k \leq m$ distinct letters j_1, \dots, j_k . If at least one of the k letters belongs to α then clearly $\mathcal{K}[\chi_{j_1} \cdots \chi_{j_k}](\alpha) = 0$. If none of them belong

to α the probability of having the given k letters in $\eta_{1,m}$ given that $\eta_{2,m} = \alpha$ is $\mathcal{K}[\chi_{j_1} \cdots \chi_{j_k}](\alpha) = \binom{n-m-k}{m-k} / \binom{n-m}{m}$. Therefore, defining λ_k as in (2.2), we have

$$\mathcal{K}\chi_{j_1} \cdots \chi_{j_k} = |\lambda_k| (1 - \chi_{j_1}) \cdots (1 - \chi_{j_k}). \quad (2.4)$$

For $k \in \{1, \dots, m\}$ let \mathcal{U}_k , be the collection of functions of the form $\varphi = \chi_{j_1} \cdots \chi_{j_k}$, with k distinct letters. Let also \mathcal{A}_k denote the linear span of $\{\mathcal{U}_0, \mathcal{U}_1, \dots, \mathcal{U}_k\}$, with \mathcal{U}_0 denoting the constant function $\varphi = 1$. Observe that by (2.4) the subspaces \mathcal{A}_k are invariant for \mathcal{K} , i.e. $\mathcal{K}\mathcal{A}_k \subset \mathcal{A}_k$. Moreover, \mathcal{A}_m coincides with $\bar{\mathcal{H}}_m$. Since \mathcal{K} is self-adjoint w.r.t. $\langle \cdot, \cdot \rangle$, to establish (2.2) it is sufficient to prove that if $\varphi \in \mathcal{A}_k$ and φ is orthogonal to \mathcal{A}_{k-1} then

$$\langle \varphi, \mathcal{K}\varphi \rangle = \lambda_k \langle \varphi, \varphi \rangle. \quad (2.5)$$

To prove (2.5) observe that $\varphi \in \mathcal{A}_k$ must be of the form $\varphi = \phi + \psi$ with $\phi \in \mathcal{A}_{k-1}$ and ψ a linear combination of functions in \mathcal{U}_k , say $\psi = \sum_i a_i \psi_i$. For each $\psi_i \in \mathcal{U}_k$ we use (2.4) to obtain $\mathcal{K}\psi_i = \tilde{\phi}_i + \lambda_k \psi_i$, with $\tilde{\phi}_i \in \mathcal{A}_{k-1}$. Setting $\tilde{\phi} = \sum_i a_i \tilde{\phi}_i$ we see that $\mathcal{K}\psi = \tilde{\phi} + \lambda_k \psi$. If φ is orthogonal to \mathcal{A}_{k-1} we then have $\langle \varphi, \mathcal{K}\varphi \rangle = \langle \varphi, \tilde{\phi} \rangle = 0$. In particular,

$$\langle \varphi, \mathcal{K}\varphi \rangle = \lambda_k \langle \varphi, \psi \rangle = \lambda_k \langle \varphi, \varphi \rangle.$$

□

We turn to a spectral gap estimate for an auxiliary block-dynamics. Suppose $m \leq n/N$ for some integer $N \geq 2$ and let I_1, \dots, I_N denote N non-overlapping m -blocks in V_n . Consider the Markov chain which picks uniformly at random one of the N blocks and, given the configuration on that block, chooses uniformly at random a compatible configuration on the complement of that block. In symbols, we are looking at the symmetric transition matrix

$$\mathcal{P}(\sigma, \xi) = \frac{1}{N} \sum_{k=1}^N \nu[\eta = \xi \mid \eta = \sigma \text{ on } I_k], \quad \sigma, \xi \in \Omega. \quad (2.6)$$

Next result is a natural extension of the estimates in Theorem 2.1 of [3] and Lemma 2.2 of [2].

Proposition 2.2. *Let μ denote the lowest non-zero eigenvalue of $1 - \mathcal{P}$. Then*

$$\mu \geq \frac{N-2}{N-1} (1 - \lambda_2) \quad (2.7)$$

Proof. For all functions $f, g : \Omega \rightarrow \mathbb{R}$ we denote by (f, g) the scalar product

$$(f, g) = \nu[fg] = \frac{1}{n!} \sum_{\eta \in \Omega} f(\eta)g(\eta). \quad (2.8)$$

Let $\pi_k : \Omega \rightarrow \Omega_m$ denote the projection $\eta \mapsto \eta_{I_k}$, the restriction to the configuration on the block I_k . We define the subspace Γ consisting of sums of mean-zero functions of a single block:

$$\Gamma = \left\{ f = \sum_{k=1}^N g_k \circ \pi_k ; \quad g_k : \Omega_m \rightarrow \mathbb{R} \text{ with } \langle g_k, 1 \rangle = 0 \quad \forall k \right\}, \quad (2.9)$$

where $\langle \cdot, \cdot \rangle$ has been defined in (2.3). We write $\mathcal{P} = \frac{1}{N} \sum_{k=1}^N \mathcal{P}_k$ with

$$\mathcal{P}_k(\sigma, \xi) = \nu[\eta = \xi \mid \pi_k \eta = \pi_k \sigma].$$

In this way $\mathcal{P}_k f$ is a function of the letters on the single block I_k only and, with a slight abuse of notation, we may identify the functions $\mathcal{P}_k f$ and $\mathcal{P}_k f \circ \pi_k$. Therefore $\mathcal{P}f \in \Gamma$ for every f with $\nu[f] = 0$. Since \mathcal{P} is self-adjoint w.r.t. (\cdot, \cdot) , $\mathcal{P}f = 0$ whenever $\nu[f] = 0$ and f is orthogonal to Γ . To prove (2.7) it is then sufficient to establish

$$(f, (1 - \mathcal{P})f) \geq \frac{N-2}{N-1} (1 - \lambda_2)(f, f), \quad f \in \Gamma. \quad (2.10)$$

Given $f \in \Gamma$, $f = \sum_k g_k \circ \pi_k$, we define $\varphi_f = \sum_k g_k$, a function on Ω_m . Using

$$(g_j \circ \pi_j, g_k \circ \pi_k) = \begin{cases} \langle g_j, \mathcal{K}g_k \rangle & k \neq j \\ \langle g_k, g_k \rangle & k = j \end{cases}$$

a simple computation shows that

$$(f, f) = \langle \varphi_f, \mathcal{K}\varphi_f \rangle + \sum_k \langle g_k, (1 - \mathcal{K})g_k \rangle. \quad (2.11)$$

Similarly, observing that

$$\mathcal{P}_k(g_j \circ \pi_j) = \begin{cases} \mathcal{K}g_j & k \neq j \\ g_k & k = j \end{cases}$$

we compute, for every k ,

$$(f, \mathcal{P}_k f) = 2\langle \varphi_f, \mathcal{K}(1 - \mathcal{K})g_k \rangle + \langle g_k, (1 - \mathcal{K})^2 g_k \rangle + \langle \varphi_f, \mathcal{K}^2 \varphi_f \rangle.$$

Averaging over k and rearranging terms we finally have

$$\begin{aligned} (f, (1 - \mathcal{P})f) &= \frac{N-2}{N} \langle \varphi_f, \mathcal{K}(1 - \mathcal{K})\varphi_f \rangle \\ &\quad + \frac{1}{N} \sum_k \langle g_k, (1 - \mathcal{K})((N-1) + \mathcal{K})g_k \rangle. \end{aligned} \quad (2.12)$$

Consider now the subspace $\mathcal{S} \subset \Gamma$ of symmetric functions:

$$\mathcal{S} = \left\{ f = \sum_{k=1}^N g \circ \pi_k ; \quad g : \Omega_m \rightarrow \mathbb{R} \text{ with } \langle g, 1 \rangle = 0 \right\}, \quad (2.13)$$

Since \mathcal{S} is invariant for \mathcal{P} , i.e. $\mathcal{P}\mathcal{S} \subset \mathcal{S}$, to prove (2.10) we may consider separately the cases $f \in \mathcal{S}$ and $f \perp \mathcal{S}$. When $f \in \mathcal{S}$, i.e. $f = \sum_k g \circ \pi_k$, we have $\varphi_f = Ng$ and rearranging terms in (2.11) and (2.12) we obtain

$$(f, f) = N(N-1) \langle g, [\mathcal{K} + \frac{1}{N-1}]g \rangle \quad (2.14)$$

$$(f, (1 - \mathcal{P})f) = (N-1)^2 \langle g, [1 - \mathcal{K}][\mathcal{K} + \frac{1}{N-1}]g \rangle \quad (2.15)$$

Since $m \leq n/2$ it is easily seen that $|\lambda_{k+1}| \leq |\lambda_k|$ for any k . Moreover, since $Nm \leq n$ we have $|\lambda_1| \leq 1/(N-1)$ so that $\mathcal{K} + \frac{1}{N-1}$ is non-negative and we may

define $\tilde{g} = [\mathcal{K} + \frac{1}{N-1}]^{\frac{1}{2}} g$. In this way we have $(f, f) = N(N-1) \langle \tilde{g}, \tilde{g} \rangle$. Clearly $\langle \tilde{g}, 1 \rangle = 0$ and $\langle \tilde{g}, \mathcal{K}\tilde{g} \rangle \leq \lambda_2 \langle \tilde{g}, \tilde{g} \rangle$, since λ_1 is negative and λ_2 is the largest positive eigenvalue (other than $\lambda_0 = 1$). From (2.14) and (2.15) we then have

$$(f, (1 - \mathcal{P})f) \geq (N-1)^2 (1 - \lambda_2) \langle \tilde{g}, \tilde{g} \rangle \geq \frac{N-2}{N-1} (1 - \lambda_2) (f, f). \quad (2.16)$$

This proves the claim for $f \in \mathcal{S}$. It remains to study the case $f \in \mathcal{S}^\perp$. For every $u \in \mathcal{S}$, $u = \sum_k u_0 \circ \pi_k$, with $\langle u_0, 1 \rangle = 0$ one has

$$(f, u) = (N-1) \langle \varphi_f, [\mathcal{K} + \frac{1}{N-1}] u_0 \rangle.$$

Thus $f \in \mathcal{S}^\perp$ implies that $[\mathcal{K} + \frac{1}{N-1}] \varphi_f$ is constant. Since $\langle \varphi_f, 1 \rangle = 0$ it follows that $[\mathcal{K} + \frac{1}{N-1}] \varphi_f = 0$. In particular, this gives

$$\frac{N-2}{N} \langle \varphi_f, \mathcal{K}(1 - \mathcal{K})\varphi_f \rangle = \frac{N-2}{N-1} \langle \varphi_f, \mathcal{K}\varphi_f \rangle. \quad (2.17)$$

The above takes care of the first two terms in the r.h.s. of (2.11) and (2.12). We now have to compare the remaining two terms. Writing $\hat{g}_k = (1 - \mathcal{K})^{\frac{1}{2}} g_k$, we observe that $\mathcal{K} \geq \lambda_1 \geq -1/(N-1)$ implies, for every k

$$\frac{1}{N} \langle \hat{g}_k, [(N-1) + \mathcal{K}] \hat{g}_k \rangle \geq \frac{N-2}{N-1} \langle \hat{g}_k, \hat{g}_k \rangle. \quad (2.18)$$

Summing over k in (2.18) we obtain the two terms in (2.11) and (2.12) and using (2.17) we see that

$$(f, (1 - \mathcal{P})f) \geq \frac{N-2}{N-1} (f, f), \quad f \in \mathcal{S}^\perp. \quad (2.19)$$

From (2.16) and (2.19) we obtain the claim (2.7) and the proof is completed. \square

2.3. Long range average-exchange block dynamics. Let ℓ and N be given integers and suppose we have exactly $n = N\ell$ vertices. We partition the vertex set V_n by means of N non-overlapping ℓ -blocks $I_{1,\ell}, \dots, I_{N,\ell}$. There are ℓ choices for such partitioning but, by symmetry, our estimates will not depend on the choice. To fix ideas, we take the ℓ -blocks $I_{k,\ell} = \{(k-1)\ell+1, (k-1)\ell+2, \dots, k\ell\}$.

The long range average-exchange block dynamics is the continuous time Markov chain obtained as follows. There is an independent rate-1 Poisson clock at each block. When block $I_{i,\ell}$ rings, a further block $I_{j,\ell}$ is chosen uniformly at random (with replacement, i.e. the choice may be $I_{i,\ell}$ itself) and an independent fair coin is tossed. If $i = j$ we do nothing, while if $i \neq j$, depending on the outcome of the coin we update the configuration over blocks $I_{i,\ell}, I_{j,\ell}$ in one of the following ways:

1. exchange the configuration on block $I_{i,\ell}$ with that on block $I_{j,\ell}$
2. choose uniformly the new configuration on $I_{i,\ell} \cup I_{j,\ell}$ among all configurations compatible with the letters outside the blocks $I_{i,\ell}, I_{j,\ell}$.

The above defined process is reversible with respect to ν and its Dirichlet form can be written as

$$\mathcal{D}(f, f) = \frac{1}{4N} \sum_{i=1}^N \sum_{j=1}^N \nu \left[(E_{i,j} f)^2 + (A_{i,j} f)^2 \right], \quad (2.20)$$

where we use the exchange gradients $E_{i,j}$ and average gradients $A_{i,j}$ defined as follows. As before, we denote π_k the projection $\eta \rightarrow \eta_{k,\ell}$ on the block $I_{k,\ell}$. For any i, j and η we define the configuration $\eta^{(i,j)}$ obtained from η after exchanging blocks $I_{i,\ell}, I_{j,\ell}$, i.e. $\pi_k \eta^{(i,j)} = \pi_k \eta$ for all $k \neq i, j$, $\pi_j \eta^{(i,j)} = \pi_i \eta$ and $\pi_i \eta^{(i,j)} = \pi_j \eta$. We agree that $\eta^{(i,i)} = \eta$. Then the exchange gradient $E_{i,j}$ is defined by

$$E_{i,j} f(\eta) = f(\eta^{(i,j)}) - f(\eta). \quad (2.21)$$

The average gradient is given by

$$A_{i,j} f(\sigma) = \sum_{\xi \in \Omega} (f(\xi) - f(\sigma)) \nu[\eta = \xi \mid \pi_k \eta = \pi_k \sigma, \forall k \neq i, j], \quad (2.22)$$

for $i \neq j$ and we agree that $A_{i,i} f = 0$.

Proposition 2.3. *Choose integers $\ell \geq 1, N \geq 2$ and set $n = N\ell$. Let $\gamma(N, \ell)$ denote the relaxation time of the long range average-exchange block dynamics. Then*

$$\gamma(N, \ell) \leq C, \quad (2.23)$$

with $C = 4 \exp(2 \sum_{k=2}^{\infty} k^{-2})$.

Proof. We first observe that $\gamma(2, \ell) \leq 4$ for all $\ell \geq 1$. Indeed, when $N = 2$ we have $A_{1,2} f = A_{2,1} f = \nu[f] - f$ so that $\mathcal{D}(f, f) \geq \frac{1}{4} \text{Var}(f)$.

We turn to the case $N \geq 3$. Using the notation (2.8) of Proposition 2.2 we write, for any $f : \Omega \rightarrow \mathbb{R}$

$$(f, (1 - \mathcal{P})f) = \frac{1}{N} \sum_{j=1}^N (f, f - \mathcal{P}_j f). \quad (2.24)$$

Note that $\mathcal{P}_j f(\eta)$ coincides with the function $\eta \rightarrow \nu[f \mid \pi_j \eta]$, the conditional expectation of f given the configuration on $I_{j,\ell}$. In particular, denoting by $\text{Var}(f \mid \pi_j \eta)$ the variance of f w.r.t. $\nu[\cdot \mid \pi_j \eta]$, we have

$$(f, f - \mathcal{P}_j f) = (f - \mathcal{P}_j f, f - \mathcal{P}_j f) = \nu [\text{Var}(f \mid \pi_j \eta)].$$

Once the ℓ variables in $I_{j,\ell}$ are frozen, the measure $\nu[\cdot \mid \pi_j \eta]$ is the uniform measure on all $(n - \ell)!$ permutations of $n - \ell$ letters over $n - \ell$ vertices. For any $\eta \in \Omega$ we may therefore estimate, by definition of $\gamma(N, \ell)$:

$$\text{Var}(f \mid \pi_j \eta) \leq \frac{\gamma(N-1, \ell)}{4(N-1)} \sum_{i \neq j} \sum_{k \neq j} \nu \left[(E_{i,k} f)^2 + (A_{i,k} f)^2 \mid \pi_j \eta \right].$$

Taking ν -expectation we then have

$$(f, f - \mathcal{P}_j f) \leq \frac{\gamma(N-1, \ell)}{4(N-1)} \sum_{i \neq j} \sum_{k \neq j} \nu \left[(E_{i,k} f)^2 + (A_{i,k} f)^2 \right].$$

Averaging over j and observing that

$$\sum_j \sum_{i \neq j} \sum_{k \neq j} \nu [(E_{i,k} f)^2 + (A_{i,k} f)^2] = 4N(N-2)\mathcal{D}(f, f),$$

we obtain

$$\frac{1}{N} \sum_{j=1}^N (f, f - \mathcal{P}_j f) \leq \gamma(N-1, \ell) \frac{N-2}{N-1} \mathcal{D}(f, f). \quad (2.25)$$

If $(f, 1) = 0$ then (2.24), (2.25) and Proposition 2.2 imply

$$(f, f) \leq \gamma(N-1, \ell) (1 - \lambda_2)^{-1} \mathcal{D}(f, f).$$

Since $\lambda_2 = \ell(\ell-1)/(n-\ell)(n-\ell-1) \leq 1/(N-1)^2$ we have

$$(1 - \lambda_2)^{-1} \leq 1 + 2/(N-1)^2 \leq \exp(2/(N-1)^2).$$

In conclusion we have shown that

$$\gamma(N, \ell) \leq \exp(2/(N-1)^2) \gamma(N-1, \ell), \quad N \geq 3. \quad (2.26)$$

Iterating this bound down to $N = 3$ we obtain the desired estimate

$$\gamma(N, \ell) \leq \exp\left(2 \sum_{k=2}^{\infty} k^{-2}\right) \gamma(2, \ell).$$

□

2.4. Extensions of the long range average-exchange block dynamics. Here we extend the dynamics defined by (2.20) to the case where n is not a multiple of ℓ . We thus consider the case $n = N\ell + m$, $1 \leq m \leq \ell - 1$. We say that a collection $D = \{I_j\}$ of subsets of V_n is an ℓ -partition if

- i) $V_n = \cup_j I_j$ and $I_j \neq I_k$ for $j \neq k$
- ii) all the I_j 's but one are ℓ -blocks
- iii) the remaining one is an m -block

Any ℓ -partition is made of N ℓ -blocks and one m -block. An ℓ -partition is called of type 1 if vertex $x = 1$ is the left end point of one of the blocks I_j . We use symbols $I_{j,\ell}$, $j = 1, \dots, N$ for the ℓ -blocks and $I_{N+1,m}$ for the m -block. Moreover, there are exactly $N+1$ ℓ -partitions of type 1, depending on the position of $I_{N+1,m}$ relative to the $I_{j,\ell}$, and we call D_k the ℓ -partition of type 1 for which $I_{N+1,m}$ is the k -th block in the partition starting from vertex 1. The elements of D_k are denoted by $I_{j,\ell}^{(k)}$ and $I_{N+1,m}^{(k)}$ ($k = 1, \dots, N+1$).

Let $\mathcal{D}_k(f, f)$ denote the Dirichlet form

$$\mathcal{D}_k(f, f) = \frac{1}{4N} \sum_{i=1}^N \sum_{j=1}^N \nu \left[(E_{i,j}^{(k)} f)^2 + (A_{i,j}^{(k)} f)^2 \right], \quad (2.27)$$

where $E_{i,j}^{(k)}$ and $A_{i,j}^{(k)}$ are defined by (2.21) and (2.22) with $I_{j,\ell}^{(k)}$ in place of $I_{j,\ell}$. Since the m -block $I_{N+1,m}^{(k)}$ is never updated, the Dirichlet form (2.27) is non-ergodic. However the average over k of \mathcal{D}_k is ergodic and we have

Proposition 2.4. *There exists $C < \infty$ such that the following holds for any $\ell \geq 1$, $N \geq 2$ and $m \leq \ell - 1$. With $n = N\ell + m$, for every function $f : \Omega \rightarrow \mathbb{R}$*

$$\text{Var}(f) \leq \frac{C}{N+1} \sum_{k=1}^{N+1} \mathcal{D}_k(f, f) \quad (2.28)$$

Proof. For $k = 1, \dots, N+1$, let J_k denote the m -block $I_{N+1,m}^{(k)}$ of the partition D_k . Let also π_k^* denote the projection $\eta \rightarrow \eta_{J_k}$. Note that by construction the m -blocks J_k have no overlap. We may then apply Proposition 2.2 defining \mathcal{P} as in (2.6) with $I_k := J_k$ and N replaced by $N+1$. In particular, for every $N \geq 2$ we have here $N+1$ blocks of length m and therefore $\lambda_2 \leq 1/N^2 \leq 1/4$. We then see that, for any f

$$\text{Var}(f) \leq \frac{N}{N-1} (1 - \lambda_2)^{-1} (f, (1 - \mathcal{P})f) \leq 3 (f, (1 - \mathcal{P})f). \quad (2.29)$$

We now observe, as in the proof of Proposition 2.3, that for each k the expression $(f, (1 - \mathcal{P}_k)f)$ can be written as $\nu[\text{Var}(f | \pi_k^*\eta)]$. Moreover, for each η , the measure $\nu(\cdot | \pi_k^*\eta)$ is uniform over the configurations of $n-m = N\ell$ given letters. For each k and η we may then apply Proposition 2.3 to the system of N blocks $I_{j,\ell}^{(k)}$, $j = 1, \dots, N$ and estimate

$$\text{Var}(f | \pi_k^*\eta) \leq C \frac{1}{4N} \sum_{i=1}^N \sum_{j=1}^N \nu \left[(E_{i,j}^{(k)} f)^2 + (A_{i,j}^{(k)} f)^2 \mid \pi_k^*\eta \right]. \quad (2.30)$$

Taking ν -expectation and averaging over k we therefore obtain

$$(f, (1 - \mathcal{P})f) \leq C \frac{1}{N+1} \sum_{k=1}^{N+1} \mathcal{D}_k(f, f). \quad (2.31)$$

The proof is completed by inserting (2.31) in (2.29). \square

2.5. Comparison estimates. The first comparison allows to go from the long range average-exchange block dynamics to a new block dynamics in which the exchange and average operations are performed on neighboring blocks only, but the exchange is speeded up by a factor N^2 .

The setting is as in Proposition 2.4 above, with \mathcal{D}_k defined by (2.27). Given the type 1 ℓ -partition D_k we agree to rename the ℓ -blocks $I_{j,\ell}^{(k)}$ so that $I_{1,\ell}^{(k)}$ comes after the m -block $I_{N+1,m}^{(k)}$, $I_{2,\ell}^{(k)}$ after $I_{1,\ell}^{(k)}$ and so on.

Lemma 2.5. *For every function $f : \Omega \rightarrow \mathbb{R}$, for every $k = 1, \dots, N+1$*

$$\mathcal{D}_k(f, f) \leq 3N^2 \sum_{i=1}^{N-1} \nu \left[(E_{i,i+1}^{(k)} f)^2 \right] + \frac{1}{2} \sum_{i=1}^{N-1} \nu \left[(A_{i,i+1}^{(k)} f)^2 \right]. \quad (2.32)$$

Proof. Let $\eta^{(i,j)}$ denote the configuration where blocks $I_{i,\ell}^{(k)}, I_{j,\ell}^{(k)}$ have been exchanged, see (2.21). Let also $T_{i,j} : \Omega \rightarrow \Omega$ denote the transformation $\eta \rightarrow \eta^{(i,j)}$. Then, assuming $i < j$ we see that

$$T_{i,j} = T_{i,i+1} T_{i+1,i+2} \cdots T_{j-2,j-1} T_{j-1,j} T_{j-2,j-1} \cdots T_{i+1,i+2} T_{i,i+1}.$$

Define $H = j - i$ and set

$$U_h = T_{i+h-1,i+h} \cdots T_{i+1,i+2} T_{i,i+1}, \quad h = 1, \dots, H.$$

Set also

$$S_h = T_{j-h-1,j-h} \cdots T_{j-2,j-1} U_H, \quad h = 1, \dots, H-1.$$

In this way, setting $S_0 = U_H$ and $U_0 = 1$ we write

$$\begin{aligned} E_{i,j}^{(k)} f(\eta) &= f(\eta^{(i,j)}) - f(\eta) = f(S_{H-1}\eta) - f(U_H\eta) + f(U_H\eta) - f(\eta) \\ &= \sum_{h=1}^{H-1} [f(T_{j-h-1,j-h} S_{h-1}\eta) - f(S_{h-1}\eta)] + \sum_{h=1}^H [f(T_{i+h-1,i+h} U_{h-1}\eta) - f(U_{h-1}\eta)] \\ &= A(f) + B(f), \quad \text{say}. \end{aligned}$$

We then estimate

$$\nu[(E_{i,j}^{(k)} f)^2] \leq 2\nu[(A(f))^2] + 2\nu[(B(f))^2].$$

We now estimate the term involving $A(f)$ as follows. By Schwarz' inequality

$$\begin{aligned} \nu[(A(f))^2] &\leq (H-1) \sum_{h=1}^{H-1} \nu[(f \circ T_{j-h-1,j-h} S_{h-1} - f \circ S_{h-1})^2] \\ &= (H-1) \sum_{h=1}^{H-1} \nu[(f \circ T_{j-h-1,j-h} - f)^2] \\ &\leq N \sum_{i=1}^{N-1} \nu[(E_{i,i+1}^{(k)} f)^2], \end{aligned}$$

where we have used the invariance of ν under all transformations S_h . The same reasoning gives the same estimate for $\nu[(B(f))^2]$. This shows that for each couple i, j we obtain the following estimate for the exchange terms:

$$\nu[(E_{i,j}^{(k)} f)^2] \leq 4N \sum_{i=1}^{N-1} \nu[(E_{i,i+1}^{(k)} f)^2]. \quad (2.33)$$

We have to estimate the average terms. To this end observe that if the symbol $\omega_{i,j}(\eta)$ is used to denote the configuration η outside the two blocks $I_{i,\ell}^{(k)}, I_{j,\ell}^{(k)}$ we may regard $A_{i,j}^{(k)} f$ as the function $A_{i,j}^{(k)} f(\eta) = \nu[f | \omega_{i,j}(\eta)] - f$. Now a simple change of variables shows that for any other block $I_{h,\ell}^{(k)}$, $h \neq i, j$:

$$\nu[f | \omega_{i,j}(\eta)] = \nu[f^{(j,h)} | \omega_{i,h}(\eta^{(j,h)})],$$

where as usual $\eta^{(j,h)}$ stands for the configuration in which the blocks $I_{j,\ell}^{(k)}, I_{h,\ell}^{(k)}$ have been exchanged and $f^{(j,h)}(\eta) = f(\eta^{(j,h)})$. We then have, using the invariance of ν under the exchange $\eta \rightarrow \eta^{(j,h)}$:

$$\begin{aligned}\nu \left[\left(A_{i,j}^{(k)} f \right)^2 \right] &= \nu \left[\left(\nu[f^{(j,h)} | \omega_{i,h}(\eta^{(j,h)})] - f \right)^2 \right] \\ &= \nu \left[\left(\nu[f^{(j,h)} | \omega_{i,h}(\eta)] - f^{(j,h)} \right)^2 \right] = \nu \left[\left(A_{i,h}^{(k)} f^{(j,h)} \right)^2 \right].\end{aligned}$$

Moreover,

$$\nu \left[\left(A_{i,h}^{(k)} f^{(j,h)} \right)^2 \right] \leq 2 \nu \left[\left(A_{i,h}^{(k)} f \right)^2 \right] + 2 \nu \left[\left(A_{i,h}^{(k)} (f^{(j,h)} - f) \right)^2 \right],$$

and Jensen's inequality implies

$$\nu \left[\left(A_{i,h}^{(k)} (f^{(j,h)} - f) \right)^2 \right] \leq \nu \left[(f^{(j,h)} - f)^2 \right].$$

The last expression is nothing but $\nu \left[\left(E_{j,h}^{(k)} f \right)^2 \right]$ and we can estimate it by (2.33). Summarizing, choosing e.g. $h = j + 1$ we have obtained

$$\nu \left[\left(A_{i,j}^{(k)} f \right)^2 \right] \leq 2 \nu \left[\left(A_{j,j+1}^{(k)} f \right)^2 \right] + 8N \sum_{l=1}^{N-1} \nu \left[(E_{l,l+1}^{(k)} f)^2 \right]. \quad (2.34)$$

Recalling the expression (2.27) for the Dirichlet form $\mathcal{D}_k(f, f)$ the bounds (2.33) and (2.34) imply the desired claim (2.32). \square

The next task is to rewrite the exchange and average contributions to the local block dynamics appearing in the r.h.s. of (2.32) in terms of the inversions $\eta \rightarrow \eta^{x,\ell}$. We start by writing each exchange term by means of three inversions. To this end we denote by z the left end vertex of $I_{i,\ell}^{(k)}$, so that $z + \ell$ is the left end vertex of $I_{i+1,\ell}^{(k)}$.

Lemma 2.6. *For every function $f : \Omega \rightarrow \mathbb{R}$, for every $k = 1, \dots, N + 1$*

$$\nu \left[(E_{i,i+1}^{(k)} f)^2 \right] \leq 3 \nu \left[(R_{z,2\ell-1} f)^2 + (R_{z+\ell,\ell-1} f)^2 + (R_{z,\ell-1} f)^2 \right]. \quad (2.35)$$

Proof. Denote by $W_{x,h} : \Omega \rightarrow \Omega$ the inversion transformation $\eta \rightarrow \eta^{x,\ell}$. It is clear that $W_{z,2\ell-1} W_{z+\ell,\ell-1} W_{z,\ell-1} \eta$ coincides with $\eta^{(i,i+1)}$. Therefore

$$E_{i,i+1}^{(k)} f(\eta) = R_{z,2\ell-1} f(W_{z+\ell,\ell-1} W_{z,\ell-1} \eta) + R_{z+\ell,\ell-1} f(W_{z,\ell-1} \eta) + R_{z,\ell-1} f(\eta).$$

The conclusion now follows from Schwarz' inequality and the invariance of ν under all transformations $W_{x,h}$. \square

Our last comparison shows how to bound each average term in (2.32) by means of inversions.

Lemma 2.7. *For every function $f : \Omega \rightarrow \mathbb{R}$, for every $k = 1, \dots, N + 1$*

$$\nu \left[(A_{i,i+1}^{(k)} f)^2 \right] \leq \frac{1}{2\ell} \sum_{x \in I_i^{(k)} \cup I_{i+1}^{(k)}} \sum_{l \leq 2\ell} \nu \left[(R_{x,l} f)^2 + (R_{x+1,l-1} f)^2 \right]. \quad (2.36)$$

Proof. The starting point is the following standard spectral gap estimate for random transpositions. Consider a system with d vertices and d letters. Let $\nabla_{x,y}$ denote the gradient associated to the transposition of the letters at vertices x and y , i.e. $\nabla_{x,y} f(\eta) = f(\eta^{x,y}) - f(\eta)$, with $\eta^{x,y}$ denoting the configuration identical to η out of x, y and such that $(\eta^{x,y})_x = \eta_y$ and $(\eta^{x,y})_y = \eta_x$. Then it is well known (see e.g. Theorem 5.1 in [2] for a simple proof) that for all $d \geq 2$, all functions f

$$\text{Var}_d(f) \leq \frac{1}{4d} \sum_{x,y} \nu_d \left[(\nabla_{x,y} f)^2 \right], \quad (2.37)$$

where ν_d is the uniform measure over the $d!$ permutations and Var_d denotes the variance w.r.t. ν_d .

We want to apply this bound to the system with $d = 2\ell$ obtained by freezing all the letters in the complement of $I_i^{(k)} \cup I_{i+1}^{(k)}$. Recall the notation $A_{i,i+1}^{(k)} f(\eta) = \nu[f | \omega_{i,i+1}(\eta)] - f$ that we have used in the proof of Lemma 2.5. We then have for each η and every $f : \Omega \rightarrow \mathbb{R}$

$$\text{Var}(f | \omega_{i,i+1}(\eta)) \leq \frac{1}{8\ell} \sum_{x,y \in I_i^{(k)} \cup I_{i+1}^{(k)}} \nu \left[(\nabla_{x,y} f)^2 | \omega_{i,i+1}(\eta) \right]. \quad (2.38)$$

Clearly, each transposition is written as the composition of two inversions: if e.g. $y = x + h$ then $\nabla_{x,x+h} f(\eta) = R_{x,h} f(\eta^{x+1,x+h-1}) + R_{x+1,h-1} f(\eta)$. Recalling that $\nu[(A_{i,i+1}^{(k)} f(\eta))^2] = \nu[\text{Var}(f | \omega_{i,i+1}(\eta))]$ the bound (2.36) is a straightforward consequence of (2.38). \square

2.6. Proof of the upper bound in Theorem 1.1. We now combine the spectral gap estimate of Proposition 2.4 with the three comparison lemmas above. Recall that we are looking for a bound of the form $\tau(n, L) \leq C(n \vee \frac{n^3}{L^3})$, i.e.

$$\text{Var}(f) \leq C(n \vee \frac{n^3}{L^3}) \mathcal{E}(f, f), \quad (2.39)$$

for arbitrary functions f , with $\mathcal{E}(f, f)$ given by

$$\mathcal{E}(f, f) = \frac{1}{2} \frac{1}{nL} \sum_{x \in V_n} \sum_{\ell \leq L} \nu[(R_{x,\ell} f)^2]. \quad (2.40)$$

For every $1 \leq \ell \leq L$ we use Proposition 2.4 and Lemma 2.5 to write

$$\text{Var}(f) \leq C N^2 E(f) + C A(f), \quad (2.41)$$

with the exchange and average terms given by

$$E(f) = \frac{1}{N+1} \sum_{k=1}^{N+1} \sum_{i=1}^{N-1} \nu \left[(E_{i,i+1}^{(k)} f)^2 \right], \quad A(f) = \frac{1}{N+1} \sum_{k=1}^{N+1} \sum_{i=1}^{N-1} \nu \left[(A_{i,i+1}^{(k)} f)^2 \right]. \quad (2.42)$$

We start by estimating the average term $A(f)$. Using Lemma 2.7 and observing that for every $x \in V_n$, on any given partition D_k the number of indices i such that $x \in I_i^{(k)} \cup I_{i+1}^{(k)}$ is at most 2 we have

$$A(f) \leq \frac{2}{\ell} \sum_{x \in V_n} \sum_{l \leq 2\ell} \nu \left[(R_{x,l} f)^2 + (R_{x+1,l-1} f)^2 \right] \leq \frac{4}{\ell} \sum_{x \in V_n} \sum_{l \leq 2\ell} \nu \left[(R_{x,l} f)^2 \right].$$

Therefore, for any $\delta > 0$ and $\ell \leq n$ such that $\delta L \leq \ell \leq \frac{1}{2}L$ we have

$$A(f) \leq 8\delta^{-1} n \mathcal{E}(f, f). \quad (2.43)$$

Summarizing, taking e.g. $\delta = 0.1$ and adjusting the value of the constant C in (2.41), we have obtained that if $\delta L \leq \ell \leq \frac{1}{2}L$, for every f

$$\text{Var}(f) \leq C N^2 E(f) + C n \mathcal{E}(f, f). \quad (2.44)$$

We turn to an estimate of the exchange terms. Observe that so far, only partitions of type 1 enter the definition of $E(f)$, see (2.42) above. However, by symmetry, we could just as well use, for any $y \in V_n$, partitions of type y defined as those ℓ -partitions $\{I_j\}$ of V_n for which the vertex y is the left end point of one of the blocks I_j . Again there are exactly $N+1$ ℓ -partitions of type y , say $D_{k,y}$, $k = 1, \dots, N+1$. Let us call $I_{j,\ell}^{(k,y)}$ the corresponding ℓ -blocks and define the associated exchange gradients $E_{i,j}^{(k,y)}$ as in (2.21) and (2.22) with $I_{j,\ell}^{(k,y)}$ in place of $I_{j,\ell}$. We may then replace $E(f)$ in (2.44) by the following average over partitions of type y , for $y = 1, \dots, \ell$:

$$\frac{1}{\ell} \sum_{y=1}^{\ell} E_y(f), \quad E_y(f) = \frac{1}{N+1} \sum_{k=1}^{N+1} \sum_{i=1}^{N-1} \nu \left[(E_{i,i+1}^{(k,y)} f)^2 \right]. \quad (2.45)$$

Now, each term $E_y(f)$ is estimated by means of Lemma 2.6. Moreover, it is easy to check that for every $x \in V_n$ and $k = 1, \dots, N+1$, there is only one vertex $y \in \{1, \dots, \ell\}$ such that x is the left end point of one of the $I_{j,\ell}^{(k,y)}$ ($j = 1, \dots, N$). It follows that

$$\frac{1}{\ell} \sum_{y=1}^{\ell} E_y(f) \leq \frac{3}{\ell} \sum_{x \in V_n} \nu \left[(R_{x,2\ell-1} f)^2 + (R_{x+\ell,\ell-1} f)^2 + (R_{x,\ell-1} f)^2 \right].$$

We want to average this expression over $\delta L \leq \ell \leq \frac{1}{2}L$. To this end choose $\delta = 0.1$ and L_0 such that the number of integers ℓ such that $\delta L \leq \ell \leq \frac{1}{2}L$ is larger than $L/4$ for every $L \geq L_0$. If $L \geq L_0$ we can therefore estimate

$$\frac{1}{\#\{\ell : \delta L \leq \ell \leq \frac{1}{2}L\}} \sum_{\delta L \leq \ell \leq \frac{1}{2}L} \frac{1}{\ell} \sum_{y=1}^{\ell} E_y(f) \leq C \frac{n}{L} \mathcal{E}(f, f),$$

for some universal constant C . Going back to (2.44) we first replace $E(f)$ by (2.45) and then make the average over $\delta L \leq \ell \leq \frac{1}{2}L$. In this way, recalling that $N = [n/\ell] \leq n/(\delta L)$ we obtain, for $L \geq L_0$ and a new universal constant C :

$$\text{Var}(f) \leq C \frac{n^3}{L^3} \mathcal{E}(f, f) + C n \mathcal{E}(f, f) \leq 2C(n \vee \frac{n^3}{L^3}) \mathcal{E}(f, f). \quad (2.46)$$

It remains to consider the case $L \leq L_0$. In this case it is sufficient to prove that $\tau(n, L) \leq C n^3$ for some universal constant C . Recall that when $L = 1$, $\tau(n, 1) \leq Cn^3$ by the well known diffusive bound for local transpositions (see e.g. [2]). On the other hand, the obvious bound

$$\mathcal{E}(f, f) \geq \frac{1}{nL_0} \sum_{x \in V_n} \nu[(R_{x,1}f)^2]$$

shows that $\tau(n, L) \leq L_0 \tau(n, 1)$, for any $L \leq L_0$. This ends the proof of the upper bound in Theorem 1.1.

2.7. Proof of the upper bound in Theorem 1.2. This is a straightforward consequence of the previous bounds on $\tau(n, L)$. Indeed, there is $C_1 < \infty$ such that $\theta^{\ell-1} \geq 1/C_1$ for any $\ell \leq (1-\theta)^{-1}$, $\theta \in (0, 1)$. Therefore, choosing an integer $L \leq (1-\theta)^{-1}$ with $L^{-1} \leq C_2(1-\theta)$, and removing all inversions with $\ell > L$ we have

$$\tau(n, \theta) \leq C_1 C_2 \tau(n, L) \leq C_1 C_2^4 (n \vee [n(1-\theta)]^3).$$

3. LOWER BOUNDS

Here we are going to find the appropriate slow modes of relaxation. As already discussed in [6, 8] the bound $O(n \vee \frac{n^3}{L^3})$ corresponds to the competition of two different phenomena: the diffusive scale $O(\frac{n^3}{L^3})$ is related to the transport of local information while the scale $O(n)$ originates from the difficulty to separate two adjacent letters during the time evolution (described in terms of conserved edges in [6]). For the sake of completeness below we derive explicitly these bounds although some of them appear already (in some cases with explicit constants) in [6].

3.1. Proof of the lower bound in Theorem 1.1.

Let the n letters be labeled by the integers from 1 to n and call ξ_x the indicator function of the event {letter n is at vertex x }. Let $g : [0, 1] \rightarrow \mathbb{R}$ be a smooth function with $\int_0^1 g(t) dt = 0$, $\int_0^1 g(t)^2 dt = 1$. Set

$$\psi(\eta) = \sum_{x \in V_n} g(x/n) \xi_x. \quad (3.1)$$

We have, as $n \rightarrow \infty$: $\nu[\psi] \rightarrow 0$ and $\nu[\psi^2] \rightarrow 1$. We may therefore estimate $\text{Var}(\psi) \geq \frac{1}{2}$ if n is large enough. The lower estimate $\tau(n, L) \geq \frac{n^3}{CL^3}$ is then a consequence of

Lemma 3.1. *There exists $C < \infty$ such that*

$$\mathcal{E}(\psi, \psi) \leq C \frac{L^3}{n^3} \int_0^1 g'(t)^2 dt \quad (3.2)$$

Proof. Write $I_{x,\ell}$ for the $\ell + 1$ -block with left end x and right end $x + \ell$. Observe that for any $x \in V_n$ and $\ell \leq L$ we have

$$\sum_{y \in I_{x,\ell}} (\xi_y^{x,\ell} - \xi_y) = 0.$$

It follows that

$$R_{x,\ell}\psi = \sum_{y \in I_{x,\ell}} (g(y/n) - g(x/n))(\xi_y^{x,\ell} - \xi_y).$$

We now expand $g(y/n) - g(x/n) = g'(x/n)(y - x)/n + O(\ell^2/n^2)$. Observing that

$$\sum_{y \in I_{x,\ell}} \nu[(\xi_y^{x,\ell} - \xi_y)^2] \leq 2(\ell + 1)/n,$$

and ignoring higher order terms we have

$$\nu[(R_{x,\ell}\psi)^2] \leq C g'(x/n)^2 \frac{\ell^3}{n^3}.$$

The claim is then obtained by averaging over $x \in V_n$ and $\ell \leq L$. \square

To prove the bound $\tau(n, L) \geq n/C$ we may proceed as follows. Let χ denote the indicator function of the event {letter 1 is adjacent to letter 2}. It is easily seen that for every $n \geq 3$ the probability of this event is $\nu[\chi] = 2/(n-1)$. The variance is given by $\text{Var}(\chi) = 4(n-2)/(n-1)^2$. The desired estimate then follows from

Lemma 3.2. *For every $n \geq 3$*

$$\mathcal{E}(\chi, \chi) \leq \frac{16}{n(n-1)}. \quad (3.3)$$

Proof. Observe that $(R_{x,\ell}\chi)^2 = \chi(1 - \chi^{x,\ell}) + \chi^{x,\ell}(1 - \chi)$. Let us analyze the contribution of $\chi(1 - \chi^{x,\ell})$ (the other term contributes the same by symmetry). If 1, 2 are adjacent in η but not in $\eta^{x,\ell}$ this implies that either $\eta_x \in \{1, 2\}$ or $\eta_{x+\ell} \in \{1, 2\}$. Therefore, for every ℓ

$$\sum_{x \in V_n} \chi(1 - \chi^{x,\ell}) \leq 4\chi.$$

Taking expectations and dividing by n this gives the desired claim. \square

3.2. Proof of the lower bound in Theorem 1.1. For the bound $\tau(n, \theta) \geq [n(1 - \theta)]^3/C$ we simply follow the computation in Lemma 3.1 to obtain

$$\mathcal{E}_\theta(\psi, \psi) \leq \frac{C(1 - \theta)}{n^3} \sum_{\ell \leq n} \theta^{\ell-1} \ell^3 \int_0^1 g'(t)^2 dt$$

The required bound now follows from the fact that $(1 - \theta)^4 \sum_{\ell \leq n} \theta^{\ell-1} \ell^3$ is uniformly bounded in $\theta \in (0, 1)$. Finally, the bound $\tau(n, \theta) \geq n/C$ is straightforward since Lemma 3.2 holds as it is with $\mathcal{E}(\chi, \chi)$ replaced by $\mathcal{E}_\theta(\chi, \chi)$.

3.3. A lower bound on the log-Sobolev constant. Here we prove the estimate (1.10). Let n be even and call ξ the indicator function of the event

$$\{\text{the first } n/2 \text{ letters occupy the first } n/2 \text{ vertices}\}.$$

Then $\nu[\xi] = \binom{n}{n/2}^{-1}$ and

$$\text{Ent}(\xi) = -\nu[\xi] \log \nu[\xi] \geq \frac{n}{C} \nu[\xi]. \quad (3.4)$$

On the other hand, only inversions $\eta \rightarrow \eta^{x, \ell}$ with $x \in \{n - L, \dots, n\} \cup \{n/2 - L, \dots, n/2\}$ can affect the value of ξ . In such cases we have $\nu[(R_{x, \ell} \xi)^2] \leq 2\nu[\xi]$ for any $\ell \leq L$. It follows that

$$\mathcal{E}(\sqrt{\xi}, \sqrt{\xi}) = \mathcal{E}(\xi, \xi) \leq \frac{CL}{n} \nu[\xi]. \quad (3.5)$$

Combining (3.4) and (3.5) we arrive at the desired estimate.

REFERENCES

- [1] S.G. Bobkov, P. Tetali, *Modified logarithmic Sobolev inequalities in discrete settings*, to appear in: J. Theoret. Probab.
- [2] P. Caputo, *Spectral gap inequalities in product spaces with conservation laws*, in: Stochastic analysis on large scale interacting systems (H. Osada and T. Funaki eds.), 53–88, Adv. Stud. Pure Math. **39**, Math. Soc. Japan, Tokyo, 2004
- [3] E. Carlen, M.C. Carvalho, M. Loss, *Determination of the spectral gap in Kac's master equation and related stochastic evolutions*. Acta Math. **191**, 1–54, 2003.
- [4] P. Diaconis, L. Saloff-Coste, *Logarithmic Sobolev inequalities for finite Markov chains*, Ann. Appl. Probab. **6**, 695–750, 1996
- [5] P. Diaconis, M. Shahshahani, *Time to reach stationarity in the Bernoulli-Laplace diffusion model*, SIAM J. Math. Anal. **18**, no. 1, 208–218, 1987.
- [6] R. Durrett, *Shuffling chromosomes*, J. Theoret. Probab. **16**, 725–750, 2003
- [7] R. Durrett, *Probability models for DNA sequence evolution*. Probability and its Applications, Springer-Verlag, New York, 2002.
- [8] R. Durrett, *Genome rearrangement: recent progress and open problems*, preprint 2003
- [9] F. Gao, J. Quastel, *Exponential decay of entropy in the random transposition and Bernoulli-Laplace models*. Ann. Appl. Probab. **13**, 1591–1600, 2003
- [10] T. Y. Lee, H. T. Yau, *Logarithmic Sobolev inequality for some models of random walks*. Ann. Probab. **26** no. 4, 1855–1873, 1998
- [11] F. Martinelli, *Relaxation times of Markov chains in statistical mechanics and combinatorial structures*. Encyclopedia of Mathematical Sciences Vol. **110**, Springer 2003
- [12] Saloff-Coste, *Lectures on finite Markov chains*. Lectures on probability theory and statistics (Saint-Flour, 1996), 301–413, LNM **1665**, Springer, Berlin, 1997.

- [13] D. Wilson, *Mixing times of Lozenge tiling and card shuffling Markov chains*, Ann. Appl. Probab. **14**, 274–325, 2004

DIP. MATEMATICA, UNIVERSITÀ DI L'AQUILA AND INFN UNITÀ DI ROMA “LA SAPIENZA”
E-mail address: nicoletta.cancrini@roma1.infn.it

DIP. MATEMATICA, UNIVERSITÀ DI ROMA TRE, L.GO S. MURIALDO 1, 00146 ROMA, ITALY
E-mail address: caputo@mat.uniroma3.it

DIP. MATEMATICA, UNIVERSITÀ DI ROMA TRE, L.GO S. MURIALDO 1, 00146 ROMA, ITALY
E-mail address: caputo@mat.uniroma3.it